

PD Dr. vet. med. habil. Wolfgang Beyer – References (Zoonoses)

University of Hohenheim, June 2016

1. Turner WC, Kausrud KL, Beyer W, Easterday WR, Barandongo ZR, Blaschke E, et al. Lethal exposure: An integrated approach to pathogen transmission via environmental reservoirs. *Sci Rep* 2016;6:27311. doi:10.1038/srep27311.
2. Ganz HH, Law C, Schmuki M, Eichenseher F, Calendar R, Loessner MJ, et al. Novel giant siphovirus from *Bacillus anthracis* features unusual genome characteristics. *PLoS ONE* 2014;9. doi:10.1371/journal.pone.0085972.
3. Terzi B von, Turnbull PCB, Bellan SE, Beyer W. Failure of Sterne- and Pasteur-like strains of *Bacillus anthracis* to replicate and survive in the urban bluebottle blow fly *Calliphora vicina* under laboratory conditions. *PLoS ONE* 2014;9:e83860. doi:10.1371/journal.pone.0083860.
4. Bellan SE, Turnbull PCB, Beyer W, Getz WM. Effects of experimental exclusion of scavengers from carcasses of anthrax-infected herbivores on *Bacillus anthracis* sporulation, survival, and distribution. *Appl Environ Microbiol* 2013;79:3756–61. doi:10.1128/AEM.00181-13.
5. Beyer W, Turnbull PCB. Co-infection of an animal with more than one genotype can occur in anthrax. *Lett Appl Microbiol* 2013;57:380–4. doi:10.1111/lam.12140.
6. Grunow R, Klee SR, Beyer W, George M, Grunow D, Barduhn A, et al. Anthrax among heroin users in Europe possibly caused by same *Bacillus anthracis* strain since 2000. *Euro Surveill* 2013;18.
7. Owen MP, Schauwers W, Hugh-Jones ME, Kiernan JA, Turnbull PCB, Beyer W. A simple, reliable M'Fadyean stain for visualizing the *Bacillus anthracis* capsule. *Journal of Microbiological Methods* 2013;92:264–9. doi:10.1016/j.mimet.2013.01.009.
8. Beyer W, Bellan S, Eberle G, Ganz HH, Getz WM, Haumacher R, et al. Distribution and molecular evolution of *Bacillus anthracis* genotypes in Namibia. *PLoS Negl Trop Dis* 2012;6:e1534. doi:10.1371/journal.pntd.0001534.
9. Welkos S, Cote CK, Hahn U, Shastak O, Jedermann J, Bozue J, et al. Humanized q-defensins (retrocyclins) enhance macrophage performance and protect mice from experimental anthrax infections. *Antimicrobial Agents and Chemotherapy* 2011;55:4238–50. doi:10.1128/AAC.00267-11.
10. Beyer W, Turnbull PCB. Anthrax in animals. *Mol Aspects Med* 2009;30:481–9. doi:10.1016/j.mam.2009.08.004.
11. Hadjinicolaou AV, Demetriou VL, Hezka J, Beyer W, Hadfield TL, Kostrikis LG. Use of molecular beacons and multi-allelic real-time PCR for detection of and discrimination between virulent *Bacillus anthracis* and other *Bacillus* isolates. *Journal of Microbiological Methods* 2009;78:45–53. doi:10.1016/j.mimet.2009.04.005.
12. Lasch P, Beyer W, Nattermann H, Stämmler M, Siegbrecht E, Grunow R, Naumann D. Identification of *Bacillus anthracis* by using matrix-assisted laser desorption ionization-time of flight mass spectrometry and artificial neural networks. *Appl Environ Microbiol* 2009;75:7229–42. doi:10.1128/AEM.00857-09.
13. Hudson MJ, Beyer W, Bohm R, Fasanella A, Garofolo G, Golinski R, et al. *Bacillus anthracis*: balancing innocent research with dual-use potential. *Int J Med Microbiol* 2008;298:345–64. doi:10.1016/j.ijmm.2007.09.007.
14. Akinyemi KO, Philipp W, Beyer W, Böhm R. In vitro antimicrobial susceptibility patterns of *Salmonella enterica* serovars and emergence of *S. typhimurium* phage type DT071 in a suspected community-associated outbreak in Lagos, Nigeria. *The Journal of Infection in Developing Countries* 2007;1.
15. Grunow R, Porsch-Özcürümez M, Splettstoesser W, Buckendahl A, Hahn U, Beyer W, et al. Monitoring of ELISA-reactive antibodies against anthrax protective antigen (PA), lethal factor (LF), and toxin-neutralising antibodies in serum of individuals vaccinated against anthrax with the PA-based UK anthrax vaccine. *Vaccine* 2007;25:3679–83. doi:10.1016/j.vaccine.2007.01.056.
16. Tomaso H, Bartling C, Al Dahouk S, Hagen RM, Scholz HC, Beyer W, Neubauer H. Growth characteristics of *Bacillus anthracis* compared to other *Bacillus* spp. on the selective nutrient media Anthrax Blood Agar® and Cereus Ident Agar®. *Systematic and Applied Microbiology* 2006;29:24–8. doi:10.1016/j.syapm.2005.05.008.
17. Nübel U, Schmidt PM, Reiß E, Bier F, Beyer W, Naumann D. Oligonucleotide microarray for identification of *Bacillus anthracis* based on intergenic transcribed spacers in ribosomal DNA. *FEMS Microbiology Letters* 2004;240:215–23. doi:10.1016/j.femsle.2004.09.042.
18. Beyer W, Bartling C, Neubauer H. Zum stand der Nachweisverfahren für *Bacillus anthracis* in klinischen und Umweltproben. *Tierärztliche Umschau* 2003;58:653–62.
19. Beyer W, Pocivalsek S, Böhm R. Polymerase chain reaction-ELISA to detect *Bacillus anthracis* from soil samples - Limitations of present published primers. *J Appl Microbiol* 1999;87:229–36. doi:10.1046/j.1365-2672.1999.00875.x.
20. Beyer W, Mukendi FM, Kimmig P, Bohm R. Suitability of repetitive-DNA-sequence-based PCR fingerprinting for characterizing epidemic isolates of *Salmonella enterica* serovar Saintpaul. *Journal of Clinical Microbiology* 1998;36:1549–54.
21. Beyer W, Bohm R. Labeling *Salmonella* live vaccine strains with the lux operon from *Vibrio fischeri* improves their detection and discrimination from wild type. *Microbiological Research* 1996;151:407–19.
22. Beyer W, Glockner P, Otto J, Bohm R. A nested PCR method for the detection of *Bacillus anthracis* in environmental samples collected from former tannery sites. *Microbiological Research* 1995;150:179–86.